Ashraf M., Berge O., Azam F. and Hentin T. 1999. Bacterial exopolysaccharides and productivity of the salt affected soils. Diversity of exopolysaccharide producing bacteria from the rhizoplane of wheat (*Triticum aestivum* L.) grown in normal and saline Pakistani soils. *Pak. J. Biol. Sci*. 2: 201-206.

Barkay T., Navon-Venezia S., Ron E.Z. and Rosenberg E. 1999. Enhancement of solubilization and biodegradation of polyaromatic hydrocarbons by the bioemulsifier Alasan. *Appl. Environ, Microbiol*. 65 (6): 2697-2702.

Beech I., Hanjagsit L., Kalaji M., Neal A. L. and Zinkevich V. 1999. Chemical and structural characterization of exopolymers produced by Pseudomonas sp. NCIMB 2021 in continuous culture. Microbiology. Vol. 145, pp. 1491-1497. ISSN 1465-2080

Bejar, V., Llamas, I., Calvo, C., Quesada, E., Characterisation of exopolysaccharides produced by 19 halophilic strains of the species Halomonas eurihalina, Journal of Biotechnology, 1998, 61, p. 135-141

Beveridge, T. J. and L. L. Graham. 1991. Surface layers of bacteria. Microbiol. Reviews 55:684-705.

Chabot S., H. L. Yu, L. De Leseleuc, D. Cloutier,M. R. Van Clasteren, M. Lessard, D. Ray, M. Lacraix, and D. Oth. 2001. Exopolysaccharides from Lactobacillus rhamnosus RW-9595M stimulate TNF, IL-6 and IL-12 in human and mouse cultured immunocompetent cells, and IFN-γ in mouse splenocytes. Lait 81:683–697.

Characklis, W.G., Escher, A.R., 1988. Microfouling in initial events. sIn: Thompson M F, Sarojini R, Nagabushanam R (eds) Marine Biodeterioration: Advanced Techniques Applicable to the Indian Ocean Oxford & IBH Publ Company Pvt. Ltd, New Delhi, Bombay, Calcutta, pp 249 260

Chemical and rheological properties of an extracellular polysaccharide produced by the cyanobacterium *Anabaena* sp. ATCC 33047. *Biotechnol. Bioengg*. 67: 283-290.

Chen J.H., Czajka. D.R., Lion L.W., Shuler M.L. and Ghiorse W.C. 1995. Trace metal mobilization in soil by bacterial polymers. *Environ. Hlth. Persp*. 103 (1):53-58.

Corpe, W. A., In *Adsorption of Microorganisms to Surfaces* (eds Bitton, G. and Marshall, K. C.), John Wiley, New York, 1980, pp.105

Crescimanno G, Iovino M, Provenzano G (1995).Influence of salinity and sodicity on soil structural and hydrolic characteristics. Soil Sci Soc AMJ 59:1701-1708.

Decho, A.W., 1990. Microbial exopolymer secretions in ocean environments: Theirrole(s) in food webs and marine processes. Oceanogr. Mar. Biol. Annu. Rev. 28,73-153.

De Philippis R., Christina M., Materassi R. and Vincenzini M. 1998. Potential of unicellular cyanobacteria from saline environments as exopolysaccharide producers. *Appl. Environ. Mocrobiol*. 64 (3): 1130-1132.

DeFilippis, V.R. and Villarreal, L.P. (2001) Virus Evolution. In: D.M. Knipe, P.M. Howley and D.E. Griffin (Eds), Fields Virology, 4th ed. Lippincott Williams & Wilkins, Philadelphia, Pa.

Ducklow, H. W., and R. Mitchell. 1979. Bacterial populations and adaptations in the mucus layers on living corals. Limnol. Oceanogr, 24: 715-725.

Emily B.Roberson and Mary K. Firestone .Relationship between desiccation & exopolysaccharide production in a soil Pseudomonas sp. Appl. Environ Microbiol.1992. April; 58(4):1284-1291.

Fernandes H.L., Lupi, F., Tome, M.M., Sa-Correia I. and Novais J.M. 1991. Rheological behaviour of the culture medium during growth of the microalga *Botryococcus braunii*. *Bioresource Technol*. **38**: 133-136.

Flecther, M. 1980. The characteristics of interfaces and their role in microbial attachment, p. 67-78. In R. C. W. Berkeley et al. [eds.], Microbial attachment to surfaces. Ellis Hoi-wood.

Garcia, C. and Hernandez, T. (1996).Influence of salinity on biological and biochemical activity of a calciothird soil.Pl soil 178 : 255-263.

Hebber P., Gueniot B., Heyraud A., Heulin T. and Rinauds M. 1992. Characterization of exopolysaccharides produced by bacteria isolated from plant roots. *Appl. Microbiol. Biotechnol*. 38: 248-253.

Helm R.F., Huang Z., Edwards D., Leeson H., Peery W. and Potts M. 2000. Structural characterization of the released polysaccharide of a desiccation tolerant *Nostoc commune* DRH-1. *J. Bacteriol*. 182(4): 974-982.

Hill, D.R., Haldun, S.L., Schere,r S. and Potts, M. 1994. Water stress protein of *Nostoc commune* (cyanobacteria) are secreted with UV-A/B absorbing pigment and associate with 1,4-ß-D-xylano hydrolase activity*. J. Biol. Chem*. 269:7726-7734.

Hill, D.R., Keenan, T.W., Helm R.F., Potts, M., Crowe, L.M. and Crowe, J.H. 1997. Extracellular Polysaccharide of *Nostoc commune* (cyanobacteria) inhibits fusion of membrane vesicles during desiccation. *J. Appl. Phycol*. 9:237-248.

Hill, D.R., Peat, A. and Potts,M. 1994. Biochemistry and structure of the glycan secreted by desiccation tolerant *Nostoc commune* (cyanobacteria). *Protoplasma* 182:126-148.

Hoa, P.T., Nair, L. and Visvanathan, C. 2003. The effect of nutrients on extracellular polymeric substance production and its influence on sludge properties. *Water S.A*. 29 (4): 437-442.

Hoagland K, Rosowski J, Gretz M, Roemaer S (1993) Diatom extracellular polymeric substances: Function, fine structure, chemistry, and physiology. J. Phycology 29:537-566.

Hoagland, K. D., Rosowski J. R., Gretz M. R. and Roemer S. C. (1993) Diatomextracellular polymeric substances: Function, fine structure, chemistry, and physiology.J. Phycol., 29, 537-566.

Hokputsa S., Hu C., Paulsen B.S. and Harding S.E. 2003. A physiochemical comparative study on extracellular carbohydrate polymers from five desert algae. *Carbohydrate Polymers*.54:27-32.

Iqbal A., Bhatti H.N., Nosheen S., Jamil A. and Malik M.A. 2002. Histochemical and physicochemical study of bacterial exopolysaccharides. *Biotechnology* **1**(1): 28-33.

Johansen J.R. 1993. Cryptogamic crusts of semi arid and arid lands of North America. *J. Phycol.* 29: 140-147.

Kurane R., Hatamochi K., Kakuno, T. and Kiyohara M. 1994. Production of a bioflocculant by *Rhodococcus erythropolis* S-1 grown on alcohols. *Biosci. Biotechnol. Biochem*. **58**: 428-429.

Kumar, C.G., Joo, H.S., Choi, J. W., Koo, Y.-M., Chang, C.-S., 2004. Purification and characterization of an extracellular polysaccharide from haloalkalophilic Bacillus sp. I-450. Enzyme and Microbial Technology 34, 673-681.

LaMourie, M.J. and P. Hoagland. 1993. Government support of marine science and technology in Japan: the case of marine electronic instrumentation. Mar. Tech. Soc. J. 27(3):21-33. [WHOI Contrib. No. 8280.

Laws, A., Gu, Y. and Marshall V. (2001) Biosynthesis, charterzation, and design of bacterial exopolysaccharides from lactic acid bacteria, Biotechnol. Adv.19:597-625.

Lowry O.H., Rosenbrough N.J., Iarr A.L. and Randall R.I. 1951. Protein measurement with Folin-Phenol reagent. *J. Biol. Chem*. 193:265-275.

Mazor G., Kidron G.J., Vonshak A. and Abeliovich A. 1996. The role of cyanobacterial exopolysaccharides in structuring desert microbial crusts. *FEMS Microbial Ecol.* 21:121-130.

Metting B. 1981. The systematics and ecology of soil algae. *Bot. Rev*. 47:195-312.

Moreno J., Vargas M.A., Madiedo J.M., Munoz J., Rivas J. and Guerrero G. 2000.

Moreno, J., Angeles Vargas, M., Madiedo, J.M., Munoz, J., Rivas, J., Guerrero, M.G., Chemical and rheological properties of an extracellular polysaccaride produced by the cyanobacterium Anabaena sp. ATCC 33047, Biotechnology and bioengineering, 2000, 67 (3), p. 283-290.

Morvan H., Gloaguen V., Vebret L., Joct F. and Hoffman L. 1997. Structure-function investigation on capsular polymers as a necessary step for new biotechnological application: The case of cyanobacterium *Mastigocladus laminosus. Pl. Physiol. Biochem*.35:671-683.

O'Brein T.P. and Mc Cully M.E. 1981. Study of plant structure. Principle and Selected Methods. Thermarcarphi Pub.. Melbourne, Australia. 276p.

Oda, M., H. Hasegawa, S. Komatsu, M. Kambe, and F. Tsuchiya (1983) Anti-tumor polysaccharide from *Lactobacillus* sp. *Agri. Biol. Chem*. 47: 1623-1625.

Painter T.J. 1993. Carbohydrate polymers in desert reclamation: The potential of microalgal biofertilizers. *Carbohydrate Polymers*. 20: 77-86.

Pigeon R. M., E. P. Cuesta, and S. E. Gilliland. 2002. Binding of free bile acids by cells of yogurt starter culture bacteria. J. Dairy Sci. 85:2705–2710.

Pearse A.G.I. 1968. Histochemistry: Theoretical and Applied. Ed.1 pp. 758.

Pearse A.G.I. 1985. Histochemistry: Theoretical and Applied. Ed.2 pp. 1055.

Pora R.J., Thompson W.A. and Kriedman D.E.C. 1989. Determination of accurate extinction coefficient and simultaneous equation for assaying chlorophyll a and b extracted with four different solvents: Verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochem. Biophys. Acta*. 975: 384-394.

Potts M.1994. Desiccation tolerance of prokaryotes. *Microbiol. Rev*. 58:755-805.

Potts M.1999. Mechanisms of desiccation tolerance in cyanobacteria. *Eur. J. Phycol*. 34 (4):319-328.

Rahman Banzegar A., Malcolm Odes J. and Rengasamy P. (1996). Soil structure degradation and mellowing of compected soil by saline sodic solution . Soil Sci Soc Am J 60: 583-588.

Rapp Telana (2012). Isolation and characterisation of genes encoding biopolymer manufacturing enzymes. *Thesis presented in partial fulfilment of the requirements of the degree Master of Science in Plant Biotechnology at Stellenbosch University .* Faculty of Agricultural Sciences Department Genetics Institute for Plant Biotechnology.

Rath, H., C. Schiller, G.J. Herndl, 1993: Ectoenzymatic activity and bacterialdynamics along a trophic gradient in the Caribbean Sea. Mar. Ecol. Prog. Ser.,10.

Rees , D. A. 1972. Shapely polysaccharides, the eighth Colworth Medal Lecture. Biochem. J. 126: 257- 273. 1976. Stereochemistry and binding behavior of carbohydrate chains, p. l-42. In W. J. Whelan [ed.], Biochemistry of carbohydrates, V. 5. Univ. Park.

Rehm Bernd H. A. (2010) Bacterial polymers: biosynthesis, modifications and applications. *Nature Reviews Microbiology* 8, 578-592.

Roe J.H. 1955. The determination of sugar in blood and spinal fluid with anthrone reagent. *J. Biol. Chem*. 212: 335-343.

Roberts IS (1995) Bacterial polysaccharides in sickness and in health. Microbiology 141:2023–2031.

Ruas-Madiedo, P., R. Tuinier, M. Kanning, and P. Zoon. 2002. Role of exopolysaccharides produced by *Lactococcus lactis* subsp. *cremoris* on the viscosity of fermented milks. Int. Dairy J. 12:689-695.

Scanlan, D. 2001. Cyanobacteria: ecology, niche adaptation and genomics. *Microbiology Today*. 28:128-130.

Scherer, S. and Potts M. 1989. Novel water stress protein from a desiccation tolerant cyanobacterium purification and partial characterization. *J. Biol. Chem*. 264: 12546-12553.

Schuster, R., Stofftransport und Reaktionskinetik bei Polysaccharid- fermentationen, Dissertation, Technischen Universitaet Muenchen, 1996.

Sheng, G. P., Yu, H. Q. and Yu, Z. (2005). Extraction of extracellular polymeric substances from the photosynthetic bacterium Rhodopseudomonas acidophila. Applied Microbiology Biotechnology. Vol. 67, pp 125–130, ISSN 1432-0614.

Shih,I.L.,Van,Y.T.,Yeh,L.C.,Lin, H.G., Chang, Y.N.,2001.Prodution of biopolymer flocculant from bacillus licheniformis and its flocculation properties.Bioresour. Technol. 78,267-272.

Shirkey, B., Kovarcik D.P., Wrigth D.J., Wilmouth G., Prickett T.F., Helm R.F., Gregory E.M. and Potts M. 2000. Active Fe containing super oxide dismutase and abundant sod mRNA in *Nostoc commune* (cyanobacteria) after years of desiccation*. J. Bacteriol*. 182:189-197.

Shivsubramanian, K. and Ganeshkumar M. 2004. Influence of vermiwash on biological productivity of Marigold.Madras Agricultural Journal. 91: 221-225.

Siddiqui S, Gupta K, Yadav A and Mangal J L (1996) . Soil salinity effect on soluble saccharides , phenol ,fatty acid ,mineral contents, and respiration rate of garlic cultivars Biologia pl 38: 611-615.

Somogyi, M. 1952. *J. Biol. Chem*. 200: 245.

Strycek T., Acreman J., Kerry A., Leppard G.G., Nermut M V and Kushner D.J. 1992 Extracellular fibril production by freshwater algae*. Microbial Ecol*. 23 (1) :53-74.

Sutherland, I. W. (1990). *Biotechnology of Exopolysaccharides*. Cambridge: Cambridge University Press.

Sutherland, I. W.(1996). A natural terrestrial biofilm. *J Ind Microbiol* 17, 281-283.

Sutherland, I. W.(1997). Microbial exopolysaccharides – structural subtleties and their consequences. *Pure Appl Chem* 69, 1911-1917.

Sutherland, I. W.(1999a). Polysaccharases for microbial polysaccharides. *Carbohydr Polym* 38, 319-328.

Sutherland, I. W.(1999b). Polysaccharases in biofilms – sources – action – consequences. In *Microbial Extracellular Polymeric Substances* , pp. 201-216. Edited by J. Wingender, T. R. Neu & H.-C. Flemming. Berlin:Springer

Sutherland, I.W. 1972. Bacterial exopolysaccharides. *Adv. Microb. Physiol.,* 8:143-212. Costerton,J.W.,Anwar,H.,&Streap,J.L.,(1992).Establishment of aging biofilms possible mechanism of bacterial resistance to antibiotic theropy.

Sutherland, J.W., A sticky business. Microbial polysaccharides: current products and future trends, Microbiology today, 2002, 29, p. 70-71.

Takeda, M., Nakano, F., Nagase, T., Iohara, K. & Koizumi, J. (1998).Isolation and chemical composition of the sheath of *Sphaerotilus natans*. *Biosci Biotechnol Biochem* 62, 1138–1143.

Tallgren, A. H., U. Airaksinen, R. von Weissenberg, H. Ojamo, J. Kuusisto,and M. Leisola. 1999. Exopolysaccharide-Producing Bacteria from Sugar Beets. *Appl. Environ. Microbiol*. 65, no. 2: 862-64.

Tuinier, R et al., (2001) Effect of structural modifications on some physical characteristics of exopolysaccharides from Lactococcus lactis, Biopolymers 59:160-166.

Van Casteran, W.H.M., Dijkema, C., Schols, H.A., Beldman, G., Voragen, A.,G.,J., Characterisation and modification of the exopolysaccharide produced by Lactococcus lactis subsp. cremoris B40, Carbohydrate Polymers, 1998, 37, p. 123-130.

Venugopal Vazhiyil (2011) Marine Polysaccharides Food Applications
CRC Press Pages 237–257 Print ISBN: 978-1-4398-1526-7.

Verhoef R. P. (2005). Structural characterisation and enzymatic degradation of exopolysaccharides involved in paper mill slime deposition. Ph.D. Thesis, Wageningen University, Netherlands.

Vicente –Garcia V.,Rios-LealE.,Calderon-Dominguze G.,Canizares-Villanueva R.O.and Olvera-Ramirez R. 2003.Detection, isolation and characterization of exopolysaccharide produced by a strain of Phormidium 94a isolated from an arid zone of Mexico. Biotechnol Bioengg. 85(3):306 -310.

Wecker, A., Modellierung der Pullulanfermentation anhand der Viskosität als indirekte Meßgröse, Fortschritt-Berichte VDI, Reihe 17, Nr. 78, Düsseldorf, VDI-Verlag, 1992.

Weigand C , Anderson G,Lingel S and Escobar D (1996). Soil salinity effets on crope growth and yield –Illustration of an analysis and mapping methodology for sugarcane. J plant physiol 148: 418-424.

Weiner R, Langille S, Quintero E (1995) Structure, function and immunochemistry of bacterial exopolysaccharides.J Ind Microbiol 15:339–345.

Whitfield C., and M.B. Perry. 1998. Isolation and purification of cell surface polysaccharides from gram-negative bacteria. *Methods in Microbiology*. 27: 249-258.

Whitton V.A. 1987. Survival and dormancy of blue green algae. *In*: Y. Henis (ed.). Survival and dormancy of microorganisms. John Wiley & Sons Inc. New York. pp. 109-167.

Wingender J., Neu T.R., Flemming, H.C., 1999. Microbial Extracellular Polymeric Substances: Characterization, Structures and Function. Springer, Berlin Heidelberg, NewYork.

Wingender J., Neu T.R., Flemming H.C., 1999. What are bacterial extracellular polymeric substances? In: Wingender, J., Neu, T.R., Flemming, H.-C. (Eds.), Microbial Extracellular Polymeric Substances: Characterization, Structure, and Function. Springer, Berlin, pp. 1–19.

Wolfaardt G.M., Lawrence J.R. and Korber D.R. 1999. Function of EPS. *In* Microbial Extracellular Polymeric Substances: Characterization, Structure and Function. Wingender J., Neu T.R. and Flemming H.C. (ed.), pp. 171-200. Springer-Verlag, New York.

Yanli M., Hongwei H., Yongsheng Y., Hongxiang O. (2012). A novel extracellular biopolymer produced from Pseudomonas fluorescens C–2: preparation, characterisation and adsorption behaviours towards nickel(II) ions in aqueous solution. *Int. J. of Materials and Structural Integrity* (6), Inderscience Enterprises Ltd pp 74-93.